

September 3rd, 1906, I found *Eurotia ceratoides* in abundance between *Camp III* (5382 m.) and *Camp IV* (5284 m.) at an altitude of about 5300 m. At 5382 m. not a single specimen was seen.

Temirlik, July 10th 1900.

Steppe region in North Tibet or rather in the open plateau-land between Akatogh and Chimen-tagh that eastwards gradually goes over into Tsaidam. Not far east of it is the lake Ghas-nor well-known from Chinese maps.

North. lat. $38^{\circ} 11'$, East. long. $90^{\circ} 19'$, Altitude 2961 m.

Mandarlik, July 13th—19th 1900.

A valley on the northern side of Chimen-tagh descending northward to Ghas-nor. There are springs and comparatively abundant vegetation.

North. lat. $37^{\circ} 47'$, East. long. $90^{\circ} 47'$, Altitude 3437 m.

Kar-yakak-sai, July 20th 1900.

A valley on the northern side of Chimen-tagh directed eastwards to the Tsaidam. Belongs to the Kwen-lun system and the regions between N. E. Tibet and Tsaidam.

North. lat. $37^{\circ} 37'$, East. long. $90^{\circ} 43'$, Altitude 3984 m.

Yapkaklik-sai, July 22nd 1900.

A valley on the N. E. side of Chimen-tagh directed to the N. E. and to Tsaidam. Belongs to the Kwen-lun system and the region between N. E. Tibet and Tsaidam.

North. lat. $37^{\circ} 32'$, East. long. $90^{\circ} 56'$, Altitude 3998 m.

Kayir, July 23rd 1900.

A valley with a brook between Chimen-tagh and Ara-tagh, N. E. Tibet.

North. lat. $37^{\circ} 26'$, East. long. $90^{\circ} 51'$, Altitude 4183 m.

Kalta-alaghan, July 24th 1900.

A mountain range in N. E. Tibet parallel to Ara-tagh and Chimen-tagh. Here *Camp XIII* is situated.

North. lat. $37^{\circ} 10'$, East. long. $90^{\circ} 43'$, Altitude 4652 m.

Ara-tagh, July 24th 1900.

Mountain range in N. E. Tibet parallel to Chimen-tagh and Kalta-alaghan. A pass in these mountains crossed July 24th, 1900, has an altitude of 4373 m.

From the neighbourhood of this pass a *Potentilla* and an *Oxytropis* were taken.

A nameless region with springs east of Kum-köl in N. E. Tibet, had an altitude of 3902 m. and was passed July 27th, 1900. *Hippuris vulgaris* was found at this place.

Kum-köl, July 28th 1900.

Freshwater lake in N. E. Tibet. At its S. E. shore is my *Camp XVI*. South of the lake and the camp an extensive field of barren sand dunes is situated.

North. lat. $37^{\circ} 17'$, East. long. $90^{\circ} 10'$, Altitude 3882 m.