

Northern *Tibet*, Camp XIII, Kalta-alaghan Mountains, 4652 m., 24th July 1900 (flow.); Mandarlik, 3437 m., medio July 1900 (flow.).

Geogr. area: Tibet.

Saussurea arenaria Maxim., in Bull. Acad. Sc. Pétersb. XXVII (1881) 490.

Northern or Inner *Tibet*, near Camp XLIV, 5127 m., 18th Aug. 1901 (flow.). Determined at Kew Herb.

Geogr. area: Kansu.

Saussurea humilis Ostf. nov. sp. (Pl. V, Fig. 4).

Caulis subnullus. Folia subrosulata linearia vel lanceolato-linearia, basi dilatata, acuta, runcinato-pinnatifida lobis parvis deflexis distantibusque acutis mucronatisve, superne \pm sparse minute glanduloso-hispida, subtus nuda, marginibus basin versus parce arachnoideo-lanatis. Capitula solitaria, rarius bina, diametro ca. 1.5 cm. Involucri bractearum ca. 3-seriatae, apice obtusae, glabrae vel apicem dorsalem versus sparse hirsutae, exteriores triangulato-obovatae, marginibus latis, nigris, interiores lanceolato-obovatae. Receptaculi setae quam achæniis breviores numerosae. Flores lilacini (?). Pappi setae 2-seriatae, exteriores quam interioribus duplo vel ultra breviores, breviter plumosae, interiores longe plumosae, ca. 10 mm. longae, basin versus fuscae, ceteroquin albæ, quam corollis breviores. Corollae tubus limbo paullo longior; antennarum appendices parce floccoso-lanatae; achænia (immatura) 2.0—2.5 mm. longa, lævia, striata angulataque.

A *S. Koslowi* C. Winkl. proxima differt foliis angustioribus glabrioribusque, apicibus involucris bractearum non reflexis; pappi setis interioribus corollae dimidium limbum attingentibus etc. A *S. Andersonii* Clarke differt receptaculis setiferis, pappi setis exterioribus breviter plumosis, achæniis lævibus etc.

Northern or Inner *Tibet*, near Camp XLIV, 5127 m., 18th Aug. 1901 (flow.).

Saussurea subulata C. B. Clarke, Comp. Ind. (1876) 226; Hook. f., Fl. Brit. India III (1882) 367; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900) 374; Hemsley, in Journ. Linn. Soc. 35 (1902) 186.

Northern *Tibet*, Mit, open valley in Kwen-lun, 4008 m., 6th Aug. 1896.

Geogr. area: Tibet, Kashmir (Himalaya) and Mongolia.

Saussurea glanduligera Schultz-Bip., in Hook. f., Fl. Brit. Ind. III (1882) 371; Hemsley, in Journ. Linn. Soc. 35 (1902) 185; Keissler, in Ann. Naturh. Hofmuseum (1907) 28; Stewart, in Bull. Torrey Bot. Club (1916) 646.

N. W. *Tibet*, the Lake Pool-tso, at the southern foot of Kwen-lun, 5077 m., 27th Sept. 1906 (flow.).

Geogr. area: Tibet, Himalaya.