

Inner *Tibet*, Camp LXVI, 4863 m., 26th Aug. 1901 (flow.); SW. Tibet, Camp CCXI, Tokchen, east of Lake Manasarovar, 4654 m., 24th July 1907 (flow. hardly begun).

*Geogr. area:* Tibet, Himalaya.

*Artemisia minor* Jacquem., ex Besser, in Bull. Soc. Nat. Moscou IX (1836) 22; Hook. f., Fl. Brit. India III (1882) 329; Hemsley, in Journ. Linn. Soc. 35 (1902) 183; Stewart, in Bull. Torrey Bot. Club (1916) 646.

Northern *Tibet*, Camp XVI, at the shore of Kum-köl, 3882 m., 28th July 1900 (flow.).

*Geogr. area:* Himalaya, Tibet. (The record from Pamir by O. Hoffmann, *Compositæ*, in O. Paulsen, Vid. Medd. Naturh. For. København [1903] 151, is wrong.)

*Artemisia Hedini* Ostf. nov. sp. (Pl. III, Fig. 1).

Sect. *Abrotanum*. Herba fragrans perennis, 10—15 cm. alta, glanduloso-pubescentis, caulibus erectis vel suberectis purpureis, ramis brevibus floriferis subadpressis. Folia sessilia, circumscriptione oblonga vel lanceolato-elliptica, subtus dense glandulosa, viridia, inferiora et media bipinnatisecta, superiora simpliciter pinnatisecta, segmentis lanceolatis vel linearibus, basin versus gradatim decrescentibus, serratis, obtusis vel mucronatis, in sicco apicibus involutis, rachi alata, irregulariter subpectinatim serrata. Capitula in racemis paniculam foliosam angustam formantibus disposita, hemisphærica, nutantia, diametro ca. 2,5 mm. Involucri squamæ late ellipticæ vel suborbiculares, glabræ vel parce glandulosæ, atræ vel atro-brunnæ dorso mediano viridi et margine late membranaceo integro vel scarioso. Corollæ extus dense glandulosæ, purpureæ; achænia juvenalia glabra.

Ex aff. *A. sacrorum* Ledeb. et *A. biennis* Willd. Ab *A. sacrorum* differt colore squamarum caulorumque, foliorum sessilium forma, indumento glanduloso, caulibus herbaceis, involucris squamis non hirsutis nec villosis. Ab *A. bienni* differt perennitate, colore squamarum, indumento glanduloso etc.

Eastern *Tibet*, Camp LXXVIII, Naktsong-tso, 4636 m., 11th Sept. 1901 (flow.).

*Senecio arnicoides* Wall. Cat. (1829) 3138, partly; Hook. f., Fl. Brit. Ind. III (1882) 350; Hemsley, in Journ. Linn. Soc. 35 (1902) 185 (var. *frigidus* Hook. f.); Stewart, in Bull. Torrey Bot. Club (1916) 646; *Ligularia arn.* Keissler, in Ann. Naturh. Hofmuseum (1907) 28.

S. W. *Tibet*, Camp CCXI, Tokchen, to the east of the lake Manasarovar, 4654 m., 24th July 1907 (flowering); height above the source of Tsangpo, 5015 m., 13th July 1907 (in bud only).

*Geogr. area:* Central and Western Himalaya; Tibet.

*Senecio goringsis* Hemsl., in Kew Bull. (1896) 212; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900) 374; *Cremanthodium g.* Hemsley, in Journ. Linn. Soc. 35 (1902) 185.

Northern *Tibet*, Camp XXV, at lake No. XVIII, 4920 m., 12th Sept. 1896.

*Geogr. area:* Tibet.