

or Inner Tibet, near Camp XLIV, 5127 m., 18th Aug. 1901 (pods nearly ripe); S. W. Tibet, Height above the source of Tsangpo, northern foot of Himalaya, 5015 m., 13th July 1907 (flowering and with young pods).

*Geogr. area:* Widely distributed in the Tibetan area, reaching into the adjoining regions of China, Mongolia and Himalaya.

*Lepidium latifolium* L., Sp. pl. ed. 1 (1753) 644; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900) 373; Hemsley, in Journ. Linn. Soc. 35 (1902) 168; Stewart, in Bull. Torrey Bot. Club (1916) 633; Fedtschenko, in Acta Hort. Petrop. XXI (1903) 285, XXIV (1904) 129, XXVIII (1909) 468.

East-Turkestan, Tatlik-bulak, south-east of Lop-nor, 1953 m., 3rd July 1900 (flowering).

N. E. Tibet, Northern slope of the south-chain of Tsaidam, 3321 m., 5th Oct. 1896.

*Geogr. area:* Temperate Eurasia, Mediterranean region including N. Africa, mostly along the sea-shores, but also in saline places in the interior; widely distributed in the inner parts of Asia.

*Lepidium cordatum* Willd., ex D. C. Regn. Veget. Syst. II (1821) 554; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900) 373; Hemsley, in Journ. Linn. Soc. 35 (1902) 168; Fedtschenko, in Acta Hort. Petrop. XXI (1903) 286, XXVIII (1909) 469.

East-Turkestan, Kamish-bulak, a spring at Bash-kurghan, 3 days journey south-east of Lop-nor, 2669 m., 5th July 1900 (flowering).

N. E. Tibet, Harato, northern slope of the south-chain of Tsaidam, 3321 m., 5th Oct. 1896.

*Geogr. area:* Siberia, East-Turkestan, Tibet.

*Dilophia salsa* Thoms., in Hook., Kew Journ. of Bot. V (1853) 20, et IV (1852) pl. 12; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900) 373; Hemsley, in Journ. Linn. Soc. 35 (1902) 168; Keissler, in Ann. Naturh. Hofmuseum Wien (1907) 22; Danguy, in Bull. Mus. d'hist. nat. (1908) 130; Fedtschenko, in Acta Hort. Petrop. XXI (1903) 286, XXIV (1904) 129, XXIV (1905) 321, XXVIII (1909) 469.

N. E. Tibet, Camp X, 5362 m., 23rd Aug. 1896.

*Geogr. area:* Tibet, Thian-shan, Kansu.

#### Fam. Fumariaceæ

(determ. by C. H. OSTENFELD).

*Corydalis Moorcroftiana* Wall. Cat. no. 1432 (1829); Hook. f. et Thoms., Fl. Ind. (1855) 266; Hemsley, in Journ. Linn. Soc. 35 (1902) 165.

S. W. Tibet, on the road between Camp CLXXXIX, Dongbo, 4598 m., and Camp CXC, Tuksun, 4596 m., Upper Tsangpo's valley; 1st July 1907 (flowering).

*Geogr. area:* Afghanistan, Tibet, Himalaya.