

Musci.

Bryum calophyllum R. Br., Suppl. App. ad it. Parryan. (1824) 196.

var. *fontanum* V. F. Brotherus, var. nov. A typo foliis laxius areolatis marginibus erectis diagnoscenda.

Northern *Tibet*, at a spring; without exact locality; ca. 4900 m., beginning of Aug. 1900.

The main species is according to communication from Dr. V. F. BROTHÉRUS before known from Tibet.

Bryum Schleicheri Schwaegr., Suppl. I., P. II, 113, 73 (1816).

Northern *Tibet*, without locality (determ. V. F. BROTHÉRUS).

Eastern *Pamir*, spring at the S. W. shore of Little Kara-kul, 3720 m., 21st July 1894 (determ. by Dr. N. BRYHN as var. *latifolia* Schimp.).

Didymodon torphaceus (Brid.) Jur., Laubmoosfl. (1882) 100.

East-Turkestan, Tatlik-bulak, 1953 m., 3rd July 1900 (determ. by Dr. V. F. BROTHÉRUS, who adds: »forma fol. cellulis lævissimis«).

Hygrohypnum palustre (Huds.) Lindb.; *Limnobium palustre*; Hemsley and Pearson, in Peterm. Mitteil. Ergänzungsbd. 28 (1900).

Northern *Tibet*, Alikhani-gol, swamp at the eastern shore of Kurluk-nor, 2780 m., 28th Oct. 1896.

Drepanocladus fluitans (L.) Warnst., in Bot. Centralbl. Beih. XIII (1903) 404.

Southern *Pamir*, Tägirmän-su, spring east of Vaktshir, 4082 m., 14th Aug. 1895, water temp. 4.5° C (determ. Dr. N. BRYHN).
